



Methodology of the Farm Carbon Calculator

And emissions factors used in reports
ending after 1 April 2025

Dr. Lizzy Parker, Dr. James Pitman, Dr. Grace
Wardell, Dr. Izzy Peters, Calum Adams and
Michael Brown

April 2025

Methodology v.3.2

Data collection spreadsheet v.1.6.3

Calculation Engine API v.1.1.1

References v.1.6.4





As a leading carbon assessment tool, The Farm Carbon Calculator is upgraded on a regular basis. This ensures our users benefit from the most recent science, new additional features and a continually improving experience. Read on to find out more.

There will be a series of updates in Spring 2025 where this methodology will be updated further. We expect changes to how the livestock section functions and as to how we calculate emissions relating to land use change. Our next scheduled interim update will be in Autumn 2025.

Table of Contents

Table of Contents	2
Glossary	3
1. About this methodology document	4
2. What's changed?	4
3. About the Farm Carbon Calculator	4
4. Standards this methodology aligns with	5
5. Independent External Review	5
6. Development cycle	6
7. Structure of the Calculator	7
8. Scope of the Calculator	7
9. Accuracy of results	8
Verification services	8
Confidence levels	9
10. References and assumptions	10
11. How do we calculate CO₂e emissions?	10
11.1. Fuels	10
Fuels and electricity	10
Travel	11
Operations	11
11.2. Materials	11
11.3. Inventory	13
11.4. Fertility & Cropping (Crops)	13
Crop emissions	13
Organic fertility emissions	14



11.5. Inputs	15
Fertilisers	15
Sprays	16
11.6. Livestock	16
Livestock	16
Animal feeds	19
Animal bedding	20
11.7. Waste	21
11.8. Distribution	21
11.9. Sequestration	21
11.10. Processing	23
12. Other Calculations we use	24
Fat and protein corrected milk (FPCM)	24
Conversions from individual GHG emissions to CO ₂ e	25
13. What farm business information do users enter?	25
14. References v.1.6.4 (1 April 2025)	26
15. Contacting us	32
16. Copyright and use	32

Glossary

AD	Anaerobic Digestion
BEIS	Department for Business, Energy and Industrial Strategy
DESNZ	Department for Energy Security and Net Zero
CH₄	Methane
CO₂	Carbon dioxide
CO₂e	Carbon dioxide equivalent
FYM	Farm Yard Manure
GHG	Greenhouse Gas
IPCC	Intergovernmental Panel on Climate Change
N₂O	Nitrous oxide
NH₃	Ammonia
PAS	Publicly Available Standard
SOM	Soil Organic Matter
SOC	Soil Organic Carbon



Document Version

Version	Date	Description
Version 1.0	August 2021	Methodology draft finalised
Version 2.0	May 2023	Methodology draft revised
Version 3.0	April 2024	Methodology draft revised
Version 3.1	October 2024	Methodology draft revised
Version 3.2	April 2025	Latest Methodology finalised

1. About this methodology document

The purpose of this document is to share details about the methodology that sits behind our Farm Carbon Calculator. With over 8000 farms actively measuring and monitoring their carbon footprint, this methodology matters. In a world grappling with the urgent task of rapidly reducing greenhouse gas emissions, we believe **transparency** in this sector is crucial.

By sharing more about how farm-related greenhouse gas emissions are measured, we hope our calculator users and the wider public will have a greater understanding about the priorities and opportunities to make positive change. We also believe transparency will help us build a greater trust and engagement with our community, and bolster feedback that will further improve our calculator.

2. What's changed?

To quickly see what has changed between this methodology and the previous version, see our separate document designed for this purpose - [found on our resources page](#).

In this process there are several items not updated and no longer available in the calculator past April 2025. Usually when this is the case, it is due to a lack of available peer reviewed data. Where gaps in knowledge exist, part of Farm Carbon Toolkit's work has been to identify, generate and fill these gaps.

3. About the Farm Carbon Calculator

The Farm Carbon Calculator is an industry-leading tool which helps farmers and growers measure, understand and take action on their carbon footprint. We are recognised as one of the UK's most trusted and fastest growing carbon tools. Recommended by the NFU and the Scottish Government, and for use in many projects, we help thousands of active users in the UK and around the world.



“Over 15 years ago, I co-created the Calculator in my spare time alongside being a grower. Created for the benefit of farmers and to help them become part of the climate solution, this ethos remains true today. With world class research behind it, over £500,000 spent on development, and thousands of users, I’m proud to see the impact this tool has had, and continues to have.” – **Jonathan Smith, Non-Exec. Director & Impact Manager**

The Calculator is part of the **Farm Carbon Toolkit**, a Community Interest Company dedicated to helping farmers and growers to transition to climate-positive practices. For over a decade, Farm Carbon Toolkit has delivered a range of practical projects, tools and services that have inspired real action on the ground. Organisations we work with include: Duchy of Cornwall, First Milk, Tesco, Yeo Valley and WWF. [Read more](#)

All users of the Farm Carbon Calculator create an account and accept Terms and Conditions which are detailed on our website: <https://farmcarbontoolkit.org.uk/terms>.

4. Standards this methodology aligns with

There is no single national or international standard which satisfactorily covers the exact requirements of a farm carbon report. Instead a range of standards are used to ensure quality and compliance.

As such we are actively moving to align with the GHG Protocol agricultural guidance, as well as land-sector based guidance from FLAG. As you will see below, the calculator makes use of the IPCC 2019 and UK GHG Inventory methodologies too.

Our tool can also be used carefully to produce carbon footprints of farm products which exceed PAS 2050:2011 requirements and which are broadly aligned with Life Cycle Analysis guidelines defined by ISO 14044 and PAS 2050 standards. PAS 2050 does not require scope 3 emissions to be included for example, and the Calculator will exceed this requirement in all use cases. In Scotland this means the tool is backed for use by the Scottish Government to fulfill the Carbon Audit requirements outlined in the Whole Farm Plan Scheme and Guidance.

If you have any questions about standards or compliance please get in touch via the details at the end of this document.

5. Independent External Review

We believe it’s important for any Carbon Calculator to be independently scrutinised and always ask or check that this is the case. We stand behind this methodology and aim to secure independent external reviews of our work on an annual basis. Our last Carbon Calculator review was completed



in February 2025 by the Carbon Trust. This reviewed the user interface, methodology, emissions factors, quality control procedures, and approach to land use change and removals against the GHG protocol, SBTi FLAG and draft LSRG, helping us to identify areas of the tool for improvement. The review highlighted key points of excellence, including:

- The tool encourages knowledge improvement around emissions reductions, with the ability to compare reports over time, there are explanations throughout the calculator and links provided in the full results breakdown to information about emissions sources.
- Users can download their reports in a range of formats (PDF, CSV, JSON).
- The quality of emissions factors - highlighting that BEIS/ DESNZ emissions factors and IPCC 2006 and 2019 emissions factors used where appropriate.
- The calculation methodologies - the IPCC 2006 and 2019 refinement have been used as the main methodological calculation within the calculator, where relevant and employing Tier 2 equations and methodologies, for example for livestock, provides UK specific emissions.
- Users can input primary data to calculate soil carbon sequestration through direct measurements and this method has been validated by a soil science academic.
- The overall layout of sections are consistent and easy to follow throughout the tool.

The review also highlighted areas requiring changes to ensure alignment with FLAG and the draft LSRG and at the time of release are:

- Improving our method to account for historical land use change and amortization of these emissions over a 20 year period
- Adding a land use tracking metric
- Separate reporting of biogenic and non-biogenic emissions
- Inclusion of leased assets
- Data quality scoring for all emissions factors
- Data entry checks to ensure the area entered does not exceed the total farm area
- Data validation checks of report start and end dates

Throughout Spring 2025, our series of updates will address these requirements, as well as bringing improved data entry options and more granularity to the tool.

6. Development cycle

The Calculator's development cycle is summarised in the figure below. The calculator is updated annually in spring though continual updates made usually in autumn - where there is a significant benefit to the end-user.

As we develop the calculator, we believe it's critical to listen to the views, requests and questions of our users to ensure we remain relevant, up to date and as user friendly as possible. We engage in a structured way which involves feedback surveys, and working groups on particular topics which are taken forward during research, development, design, and testing phases.




Figure: Farm Carbon Calculator's continual calculator development cycle

7. Structure of the Calculator

The Calculator is split into ten sections, each subdivided into various input fields and produces a report, which can be viewed or exported in a number of ways. Users enter data based on the following guidance:

- What is relevant to their business only
- Looking over the previous 12 months from a single point in time
- Including capital items like machinery and buildings that were purchased during the reporting period within the Inventory section.

8. Scope of the Calculator

The Calculator is foremost a whole farm carbon footprinting tool but can also be used to produce a footprint for each product being produced on a farm - wheat, milk, potatoes for example.

The boundary of the footprint is decided by the user and can be one of three options:

1. **To farm gate only** – i.e. no transport of produce
2. **Farm and distribution** – i.e. including transport to the customer
3. **Farm and distribution through to final customer** – i.e. including processing, and transport to the end customer's doorstep



The Calculator can also be used to footprint other businesses such as processors, distributors or wholesalers, or be used to deliver footprints of farms on the above basis as a service. These are paid services, see [our services page](#) for details.

The Calculator covers Scopes 1, 2 and 3 in its calculations:

Scope 1	Also known as direct emissions , these are emissions that are owned or controlled by the company such as tractors, farm machinery, gas for heating and from change of land use. Additional emissions arise from N ₂ O released as a consequence of manure storage and application.
Scope 2	These are associated with emissions resulting from the generation of purchased electricity used on the farm.
Scope 3	Also known as indirect emissions , associated with the production, processing and distribution of inputs into the farming system. For example, fertilisers and the emissions that occurred in the manufacture of machinery, building materials and other farm infrastructure.
Out of scopes	These are emissions associated with the combustion of biofuels, wood or crop biomass.

Users are encouraged to be as comprehensive as possible with the data they submit for their calculation, as this gives more assurance in terms of the reliability of the results.

All GHG fluxes are reported in the standard tonnes of CO₂e. In the final report, a breakdown of fluxes from carbon dioxide (CO₂), methane (CH₄) and nitrous oxide (N₂O) in tonnes of CO₂e is given, as well as a breakdown of fluxes by scope.

9. Accuracy of results

The accuracy of a carbon footprint report is dependent on a number of factors, including:

- Accuracy of emissions factors;
- Whether a factor is based on actual or proxy values;
- Accuracy of both data collection and data input by the user;
- Level of completeness by the user.

Verification services

At present we do not offer verification of carbon reports for standard users of the Calculator as doing so would require a detailed audit process. We can validate your report – which usually involves a desk-based assessment of its completeness and accuracy, before checking and



communicating the results to you. To enquire about this service and how it can help you - [get in touch](#).

As part of our consultancy service our advisors support farms and companies with enhanced footprint calculations where we also verify the inputs - which provides a level of independent auditing our clients need. We do not currently provide this service to a Third Party verification standard e.g. an ISO standard. To find out more about how we can help you see: [Our Services - Farm Carbon Toolkit](#)

Proxy and actual data

Some emissions factors are calculated based on actual data (e.g. litres of red diesel used), and some are based on proxy data (e.g. carbon sequestration of hedgerows). This depends on the availability of reference data for a particular item, and how practical it is for the user to provide data. Some items offer a choice between approaches depending on what information the user has access to – e.g. when tracking the emissions of a car, users have the option to fill actual data or proxy data. A user can either input fuel usage for their car directly if the fuel volume has been logged, or alternatively if the volume has not been recorded users can input the mileage driven by the user. The direct fuel usage provides a more accurate assessment of emissions, whereas the mileage provides a proxy value.

Users will not always have access to the equipment to directly measure GHG fluxes on their farms and so even where a user chooses the “actual” option to input data, the report for a farm or product is still an indirect assessment of its carbon footprint.

Users can, however, input direct measurements of soil organic matter (SOM) or soil organic carbon (SOC) which can be used as direct measures of GHG fluxes from soils. Indirect estimations of sequestration and land use related emissions can also be selected in the Calculator where SOM and SOC sampling is not available. If you are seeking to enter the voluntary carbon market, you should check the requirements of any scheme (more guidance here <https://farmcarbontoolkit.org.uk/toolkit-page/getting-paid-for-carbon/>).

Confidence levels

In the full results of carbon reports we provide a confidence level column. This ranges from 1 to 3, where results with 3 are those in which we have the most confidence in results. This scale is created by us through an understanding of the accuracy of the emissions/sequestration factors, as well as the likely limitations of user accuracy. For example for emissions from diesel we score this as a 3, because the emissions factors are accurate and we would expect users to also have detailed information on their usage. Conversely, emissions from livestock are scored 1 because whilst users will likely have detailed input data, the inherently variable emissions from biological systems like livestock limits the level of certainty we can have in these results.

10. References and assumptions

The majority of the emission and sequestration factors that underpin the Farm Carbon Calculator are found within peer-reviewed scientific papers and official government sources, we are transparent about these sources. These references and factors are reviewed and updated annually as part of our update cycle. A full list of current references and assumptions is provided on our website here: [References \(https://calculator.farmcarbontoolkit.org.uk/references-0\)](https://calculator.farmcarbontoolkit.org.uk/references-0) and at the end of this document.

For ease of use, our calculator is divided into the following data input categories and in subsequent sections of this document, we cover the methodology and emission factors used in each:

- Fuels
 - Materials
 - Inventory
 - Fertility & Cropping (Crops)
 - Inputs (agro-chemicals)
 - Livestock
 - Waste disposal
 - Distribution
 - Land Use & sequestration
 - Processing
-
- Waste (legacy) - This section remains available in older reports created prior to April 2024 for backwards compatibility. This was superseded by 'Waste disposal' above.

11. How do we calculate CO₂e emissions?

11.1. Fuels

Emissions from the use of fuels, electricity, travelling and contractors. These include scope 1 (direct), scope 2 (indirect emissions from purchased energy) and scope 3 (indirect – such as processing and transport) emissions, including 'well-to-tank' emissions factors.

Fuels and electricity

All of the items in liquid fuels, electricity, gas fuels, heat & steam, solid fuels, accommodation, public transport and contractors are derived from DEZNS UK GHG inventory conversion factors (107). The exceptions are:

Section	Item	Reference	Notes
Liquid fuels	AdBlue	69	



Electricity & Gas Fuels	Electricity/Gas exported to the grid	61	GHG protocol agricultural guidance on how electricity export is recorded
Electricity	Tariff with known carbon footprint	N/A	To enable users to input a known carbon footprint of an electricity supplier. Simply direct input of a CO ₂ e figure.
Gas Fuels	Biogas for Off grid	38	Accounting for gas burnt on site but generated from AD plants.
Deliveries	Known carbon footprint	N/A	To enable users to input a known carbon footprint of deliveries. Simply direct input of a CO ₂ e figure.
Operations	My Operations	37	Emissions factors are based on average fuel usage for the operation and the UK GHG inventory conversion factors.
	Contractors Operations (C.O)		

Travel

All data is from the UK GHG inventory conversion factors (107) and includes all scope 3 emissions, including ‘well-to-tank’ emissions factors. The ‘miles per gallon’ function for cars is calculated as a function of miles travelled divided by miles per gallon, to calculate gallons of fuel used. The emissions factor for petrol or diesel in litres is then multiplied by the conversion factor for litres to gallons.

Operations

Users can enter various farm activities under this header based on whether they have carried out the operations themselves or have a contractor undertaking them. This enables the operations to be taken into account if fuel usage is unknown. **If fuel usage is known, this can be entered under Liquid fuels> Diesel> Red Diesel and users should not double count it here.** Field operation data draws from the AHDB’s HGCA Calculator (37), multiplied by the diesel emissions factor (scopes 1 & 3) from the UK GHG inventory conversion factors (107). For contracted emissions these will all fall under scope 3 emissions, whereas your own field operations entered this way will be split between scope 1 and scope 3. Additional calculations made for the following options under field operations:

Section	Item	Notes
Hay baling	Small rectangular	Assumes 250 bales/ha
	Large round	Assumes 15 bales/ha
	Heston	Assumes 7.5 bales/ha

11.2. Materials

The embodied energy in a range of materials that may be used on farms, including aggregates, metals, wood and plastics. These are all Scope 3 emissions.



Emissions factors are drawn from the Inventory of Carbon and Energy (ICE) database, either version 2.0 (2), [version 3.0](#) (2a) or version 4.0 (108). Priority is given where possible to the latest version 4.0, then 3.0, and lastly 2.0. A range of metrics are used, including tonnes, kg, m² and m³. The exceptions from this source are listed below, with some being derived from factors in the ICE database, and not drawn directly from ICE:

Section	Item	Reference	Notes
Aggregates	Recycled asphalt	60	Allows the asphalt factor to be adjusted for recycled content
Various	Plastics	107	The plastic emissions factors are taken from the UK GHG inventory conversion factors database.
Fencing	Complete fencing options & components	108 & Calc	Calculating the posts and wire used in common fencing options, multiplied by emissions factors from the Inventory of Carbon and Energy.
Vineyard trellising	Vineyard trellises	107 & 108	Calculations for trellises based on the materials used
Consumables Packaging	Various	107 & Calc	The emissions factors are calculated based on average weight of the item and material used
Consumables agriculture	Bale wrap	107 & Cal	Factors by the bale provided based on average weight of material used
Horticultural materials	Netting	107 & Calc	Factors for netting based upon material usage
Horticultural constructions	Poly tunnels	108 & Calc	Factor calculations based on material usage for standard polytunnel constructions
Surfacing	Surfaces, subbase, decking, etc.	107 & Calc	Factor for materials from ICE, and area emissions factor based on calculation for surface requirements.
Computers	Laptops & Desktops	109	Proxy emissions factors for embodied energy in computers from IDEMAT
Water	Mains water & sewage	107	Scope 3 emissions for water supply and disposal
Water	Non-mains	N/A	Figure simply recorded as water use. No emissions specifically – any fuel or electricity used in pumping or treatment will be picked up under Fuels.
Cleaning products, detergents, etc.	Various	103	Product specific emissions. "Product not listed" options in each section are the average of specific product emissions included in that section.



11.3. Inventory

This section covers the embodied energy in larger items like machinery and buildings (capital items). The GHG protocol guidance advises that all capital items are accounted for ‘up-front’ in the year of emissions. This can mean spikes in carbon footprints associated with inventory, and therefore by entering capital items through this tab you have the ability to easily separate out these emissions from the rest of your footprint. We also offer the option to account for capital items over a period of 10 years, although this is not GHG protocol compliant. This is a similar principle to financial accounting, as the capital item emissions are depreciated over 10 years, so 10% of emissions are apportioned each year.

Most of the emissions factors for inventory items are again derived from the Inventory of Carbon and Energy, version 2.0 (2), [3.0](#) (2a) or 4.0 (108). It is also possible to create “custom” projects and group together any items from the “Materials” section to be treated as capital items. The other data sources are:

Section	Item	Reference	Notes
Vehicles	Cars	91	Values from the Average of all GM vehicles produced and used in the 10 year life-cycle.
Farm machinery	Tractor, harvesters, etc	3	Based on horsepower of machine – a proxy for emissions
Agricultural buildings		108 & calc	This calculation is based on a standard agricultural portal building constructed of concrete floor, steel frame, roof sheets and timber slat walls. Based on a per m2 calculation.

11.4. Fertility & Cropping (Crops)

This section covers the carbon dioxide and nitrous oxide emissions from fertility and biomass inputs to cropping systems (from varying organic fertility sources).

Crop emissions

Emissions from crops are specifically worked out from the amount of crop (fresh yield) that results in crop residues. Crop residues contribute nitrogenous material to the soil, some of which goes through denitrification to N₂O. To give a more accurate representation of how much crop residue has been left in the field, multiple levels of crop residue management practices are available for input into the calculator. For perennial crops such as soft fruits, top fruits, biomass crops and green manures, temporary grasses and cut forages, the renewal rate (i.e. the frequency at which plants are removed and replaced with new seeds, seedlings or rootstocks) of the plant is included in the calculation so it does not overestimate the amount of crop residues.



The methodology used is that of the IPCC 2019 (94), using emissions factors specific to the UK from the UK GHG Inventory and its annexes (111a & 111b) with reference to the GHG protocol agricultural guidance (61).

Some crops (for example Christmas Tree crops) have been included for data capture only and do not currently have an emissions factor associated with them.

Organic fertility emissions

The application of organic fertility sources will result in N₂O emissions as the nitrogen content of the product undergoes de/nitrification by soil bacteria, which is then volatilised into NH₃ and NO_x, and is leached or runs off from where it is applied. To calculate these emissions we use the IPCC methodology for N₂O emissions from managed soils (94), with nitrogen content data pulled from the RB209 (96b), direct N₂O emissions factors drawn from analyses of UK agricultural soils (51), and indirect emissions factors from the UK GHG inventory (111) and the IPCC (94). These sources allow us to calculate the emissions coming from organic fertility applied to grassland or arable soils during different periods of the year, with separation of different products (from separated slurry components to chemically treated paper crumbs). Conventional application will often have the highest emissions associated with it, and therefore we have included options for alternative application approaches and for post-spreading incorporation based upon in-field research (114).

Section	Item	Reference	Notes
Crops	Agricultural	111b & 94	IPCC methodology, and factors from UK GHG Inventory
	Horticultural	111b & 94	IPCC methodology, and factors from UK GHG Inventory
Market Garden		111b & 94	IPCC methodology, and factors from UK GHG Inventory, item entry in kg or per unit scale
Biomass Crops	Willow coppice	111b & 94	IPCC methodology, and factors from UK GHG Inventory
	Poplar coppice		
	Miscanthus		
	Hemp		
	Switchgrass		
Green manures, temporary grasses and cut forages	All leguminous and non-leguminous green manures and managed perennial grasses	111b & 94	N ₂ O emissions as part of the N fixation process. IPCC methodology, and UK specific N fixation rates. Note that this does not take account of any carbon sequestration – this is covered under soils in the sequestration tab. Users can enter different crop

			management regimes. Unmanaged grassland should not be entered here.
Organic fertility sources		51, 94, 96, 111, & 114	Emissions are calculated as per the IPCC methodology or N ₂ O from managed soils, using UK data sources for N content application approach.
Anaerobic digestion	Running an AD plant	7 & 38	Average emissions of various processes in running an AD plant, including CO ₂ and CH ₄ emissions. Based on tonnes of biowaste input.
Lime & Mineral fertilisers	Lime, rock phosphate, rock potash, K fertiliser, Gypsum	3 & 111c	Emissions from processing of lime and mineral fertilisers
	Phosphoric acid	109	Emissions associated with production of the amendment
	Potassium sulphate	90	
	Sulfuric acid	109	
Plant raising media		16	Average of emissions for all common plant raising media used in horticulture using the LCA approach

11.5. Inputs

The GHG emissions associated with energy input in the production of agro chemicals and, in the case of fertilisers, the N₂O emissions resulting from their application to UK soils.

Fertilisers

This is split into two sections: one is for generic fertilisers, such as Ammonium Nitrate (Product with 33.5% N) or Urea. These are derived in two parts; the manufacturing emissions from Brentrup *et al.* 2018 (48), and the application emissions from IPCC chapter 11, N₂O emissions from managed soils (94). These fertilisers require the user to specify the country of origin which should be provided on the invoice or labelling (and has a big effect on the carbon footprint of the product).

The second section is for specific solid or liquid fertilisers, including those manufactured by Yara, CF, Origin and Mole Valley Farmers. These are derived from either communication of the recipe and production methods directly from the manufacturer and then calculation using the generic fertiliser values (CF and Mole Avon) or based on verified and certified carbon footprints of those products (47, 48, 49).

The user input figures are based on tonnes or litres of product used.

Two further functions enable users to enter:

1. A specific blend of fertiliser, based on known % of N:P:K, multiplied by tonnes of product used



2. A specific known footprint of a fertiliser, using kg of CO₂e per kg of product, multiplied by tonnes of product used

Overall GHG emissions for fertilisers are based on four processes, and measured in tonnes CO₂e:

- Production emissions to factory/plant gate
- Direct N₂O emissions to soil
- Indirect NH₃ and NO_x losses (to leaching and volatilization)
- Emissions from urea hydrolysis (applies to Urea products only)

All calculations are based on IPCC methodology. The emissions factors for in field emissions are based on [MIN-NO project](#) findings (47), which are UK specific, and considered an improvement on IPCC methodology because they are more accurate.

Application is assumed to be by broadcast or application of solution. Nitrogen inhibitors are not accounted for.

Sprays

Sprays can be entered as either “generic” or “actual” depending on whether the product in question is listed in our database. Both rely on the same underlying emissions factors for fungicides, growth regulators, herbicides, insecticides, molluscicides or adjuvants (18, 40) multiplied by the concentration of active ingredient used. For “actual” sprays, we have a database of over 6000+ specific branded sprays and their active ingredient content taken from the [UK pesticides register](#).

11.6. Livestock

This section covers nitrous oxide and methane emissions from animals' enteric fermentation and manures, and emissions from imported feeds and bedding.

Livestock

Livestock emissions are complex and are based on IPCC calculation methodologies. There are several variables which require user input:

- Category of livestock, by species, age, use and live weight
- Numbers of livestock, on average, per year – both for the current year and the previous year
- Manure handling – the percentage (on an annual basis) of manures handled as slurry, FYM, daily spread, or in-field.
- Adjustments for dairy cattle (based on annual milk yield) and beef cattle (based on average liveweight).

A full list of livestock categories used in the Calculator and their default liveweight is available below:

Category	Category description	Live weight (kg)	
Dairy cattle	Dairy cows	Lactating, "dry" or in-calf dairy cows	685
	Dairy heifers	First pregnancy or first lactation dairy cattle under 3 years of age	466
	Dairy replacements (1+ years)	1-3 year old female cattle to join the dairy herd who are not in-calf or lactating	466
	Calves (under 1 year)	Cattle under 1 year old	185
	Dairy beef (1+ years)	Dairy breeds not lactating but fattened for beef (over 1 year old)	550
	Bulls for breeding	Dairy or beef breeding bulls	900
Beef cattle	Calves (under 1 year)	Cattle under 1 year old (male or female)	185
	Beef cattle	12-18 months cattle for finishing (male or female)	385
	Beef finishing heifers	18-30 months heifers for slaughter	600
	Beef suckler cows	Lactating, "dry" or in-calf beef suckler cows	550
	Bulls for breeding	Dairy or beef breeding bulls	900
Beef cattle (continued)	Finishing bulls (beef)	Bull for beef 12+ months (e.g. cereal-fed)	900
	Beef replacement heifers	First pregnancy or first lactation beef suckler cows under 3 years of age	400
	Beef finishing steers	12-24 months steers for slaughter	600
Pigs	Adult sows	Sows (including sows in pig, sows being suckled and dry sows being kept for further breeding)	185
	Breeding gilts (female)	Gilts (including gilts in pig and gilts not yet in pig)	110
	Adult boars	Boars for service	200
	Piglets	Fattening swine under 20 kg	5
	Weaner pigs (under 20kg)	Fattening swine under 20 kg	15
	Weaner pigs (over 20kg)	Fattening swine 20-80 kg	30
	Finishing pig (porker)	Fattening swine 20-80 kg	77
	Finishing pig (cutter)	Fattening swine 80+ kg	88
	Bacon pigs	Fattening swine 80+ kg	94
	Barren sows for finishing	Barren sows for fattening >80kg	185

Category		Category description	Live weight (kg)
Sheep	Ewes	Adult ewes	70
	Replacement ewes	Shearling or replacement ewes (1+ years)	60
	Rams or tups	Adult rams or tups	110
	Lambs	Young sheep under 1 year	25
Poultry		Chickens – layers	2.25
		Chickens – broilers	2.25
		Chickens - pullets	2
		Breeding stock (all poultry)	0.045
		Ducks	3.25
		Turkeys	13.2
		Geese	7.5
		Pheasants	1.2
Other livestock		Goats	50
		Horses	450
		Deer (all)	60
		Llamas	60
		Alpacas	110

Please see notes in our [Livestock Wizard](#) or [data collection spreadsheet](#) for guidance on completing this section of the Calculator (including how to estimate average head of animals in each category over the 12 month reporting period).

Emissions factors that the calculations are based on are determined by UK GHG inventory and its annexes (111) and IPCC methodology 2019 (94). Since the sex and age of the animal affects their metabolism, and therefore their enteric methane (CH₄) emissions and excretion rate, livestock are separated by these characteristics in order to improve the estimates of GHG emissions, which are inherently variable. Lactation and pregnancy also alter an animal's GHG emissions so livestock are also separated based on this trait.

Within the Calculator, it is possible to simply enter only the average head of livestock in each applicable category for the most basic estimation of GHG emissions. In this case, where no liveweight is entered, a default liveweight is used (for categories of growing livestock, e.g. calves, this



is a midpoint weight within the age-range, to take account of growth across the 12 month reporting period). These default values can be found in the table above.

For a more comprehensive estimation of GHG emissions, we recommend creating multiple entries for each category of animal and entering your own liveweights as this will give a more accurate and representative estimate of GHG emissions. This can be further increased by inputting information on dry matter intake (DMI) per head per year if this is known. Our new [Livestock Wizard](#) is designed to make calculating these averages easier for the user for a group of livestock of your choice.

A Tier 2 (UK-specific) methodology is employed to calculate livestock GHG emissions for cattle, sheep, and pigs. Poultry calculations undergo a Tier 2 calculation but with a zero value for enteric emissions while goats, horses and deer are treated with a Tier 1 (international) methodology.

In this way, the Calculator's Livestock section is customisable for a range of livestock production systems, whilst relying on the generic livestock categories underpinned by the IPCC and UK GHG Inventory guidance.

Unfortunately, the IPCC guidelines do not currently incorporate a comprehensive GWP* methodology and there is no consensus on how this methodology would be used. Our teams are working in this area and monitoring guidance as it develops but this does not yet form part of this current methodology.

Animal feeds

These indirect emissions are very important to assess the holistic carbon impacts of livestock production. If feed has been grown on-farm, users can enter it under 'Feed by-products of on-farm cropping'. Users should still enter the relevant crop / yield information under the crops section with the appropriate residue management option.

The Calculator primarily uses data from the '[GFLI database](#)' (105). Some further data for non-organic feed is obtained from '[GHG emissions from food](#)' (17), along with all the data for organic feeds.

Some emissions factors for feed blends and supplements have been calculated, based on the known constituents of certain blends. This research has been undertaken by Farm Carbon Calculator, based on discussions with feed companies. Using the constituent parts, and data from the GFLI database, the footprint of certain blends has been calculated.

Section	Item	Reference	Notes
Generic Feed blends	16% CP Dairy blend	105 & Calc	Barley/Wheat/Maize [30%], Sugar Beet Pulp [15], Soybean Meal [12], Rapeseed Meal [15], Distillers' Grains [10], Soya Hulls [10], Molasses [6], Minerals & Vitamins [2]

	18% CP Dairy blend	105 & Calc	Barley/Wheat/Maize [28%], Sugar Beet Pulp [14], Soybean Meal [18], Rapeseed Meal [12], Distillers' Grains [10], Soya Hulls [10], Molasses [6], Minerals & Vitamins [2]
	21% CP Dairy blend	105 & Calc	Barley/Wheat/Maize [25%], Sugar Beet Pulp [13], Soybean Meal [25], Rapeseed Meal [14], Distillers' Grains [5], Soya Hulls [10], Molasses [6], Minerals & Vitamins [2]
	24% CP Dairy compound	105 & Calc	Maize/Wheat/Barley [27%], Sugar Beet Pulp [15], Soybean Meal [30], Rapeseed Meal [15], Distillers' Grains [10], Soya Hulls [10], Protected Proteins [10], Molasses [5], Minerals & Vitamins [3]
	18% Fibre blend	105 & Calc	Sugar Beet Pulp [30%], Soya Hulls [30], Wheat Bran [15], Rapeseed Meal [12], Distillers' Grains [5], Molasses [6], Minerals & Vitamins [2]
	18% starch compound	105 & Calc	Maize/Wheat/Barley [45%], Sugar Beet Pulp [15], Soybean Meal [14], Rapeseed Meal [13], Distillers' Grains [5], Molasses [6], Minerals & Vitamins [2]
Calf rearing	Whole milk powder	68	Analysis of production within the research paper.
	Milk replacement powders (all)	105, 67 & 68	Formulations of milk replacement powders taken from 67 and relevant emissions factors applied to constituent parts based on information in 105 and 68
	Calf rearing pellets	105 & 67	Formulations of milk replacement powders taken from 67 and relevant emissions factors applied to constituent parts based on information in 105.
Supplements	Novapro	72	Estimate of emissions associated with constituents of Novapro (factor to be reviewed upon acceptance of product into GFLI database)

Animal bedding

Animal bedding materials emissions factors are taken from the Inventory of Carbon and Energy (ICE) database v4.0 (108) and from the GHG emissions of various straw (17) with users entering tonnes of product for an annual reporting period. If bedding has derived from on-farm production, users can enter it under 'Bedding by-products of on-farm cropping'. Users should still enter the relevant crop/ yield information under the crops section with the appropriate residue management option.



11.7. Waste

This section covers emissions from landfill, and the savings from recycling and composting materials. Users enter data on their annual outputs of waste and recycling from a range of specific categories of materials.

Emissions factors for all waste disposal emissions, including landfill, recycling, combustion and composting come from the UK GHG inventory conversion factors (107).

11.8. Distribution

For businesses that have distribution beyond the farm gate within the scope of their report, this section calculates the emissions from distributing and refrigerating food products.

Users can enter actual data on fuel used per year on distribution. If they don't have this data they can use proxy data based on three variables – delivery distance per journey, weight carried per journey, and number of journeys per year.

All the emissions factors are derived from the UK GHG inventory conversion factors (107). Average values are used, and for road haulage this is based on 50% laden lorries (on a round trip).

Users are encouraged to carefully map and describe the scope of the study, and at what point the responsibility for food transport is passed on to the next actor in the supply chain. This will be different for every business, and may range from farm gate all the way through to the customer's house.

Refrigeration emissions are calculated from refrigerant losses from food storage on the farm (or in packhouses/warehouses/food processing). This is calculated by using the GHG protocol worksheet (12), an online tool to calculate the accurate emissions from refrigerant gases, per year. The figure from the spreadsheet can then be entered directly into the Calculator by the user.

Users are reminded not to double count any data entered in the Fuels section in Distribution (and vice versa).

11.9. Sequestration

This section calculates carbon sequestered by perennial plants and soils on the farm.

Data sources: All of the sequestration factors are proxy figures, except for actual Soil Organic Matter (SOM) or Soil Organic Carbon (SOC) measurements. A range of sources are used in this section:

Section	Item	Reference	Notes
Soils	Soil Organic Matter	79	Based on actual SOM and/ or SOC from soil samples, users enter data on field size, depth of measurement, bulk density and SOM/SOC results over a given time period.

	Soil Organic Carbon		This is converted into changes in volume of soil organic carbon and therefore the amount of carbon sequestered (or emitted) as per IPCC methodology.
	Carbon stocks		A log of baseline soil carbon stocks in fields. These results do not impact on the overall carbon balance, they are therefore just for reference.
Woodland	Detailed analysis	104	Users input the species, age range and area of woodland. Assumptions of average yield class, average spacing, and no thinning is applied. This is the recommended approach.
	Mixed, coniferous and broadleaf	104	Average values per hectare of types of woodland, over a 200 year average.
	In field trees	104	A per m ² value based on average sequestration rates for deciduous woodland.
Hedgerows	Managed (generic)	22, 25, 99, & 101	Based on the length and width of managed hedges – i.e. those cut on a regular basis. Sequestration factors based on averages from peer reviewed studies.
	Managed hedgerow under 15 years old	87, 88, 89	Based on the length and width of managed hedges – gives age-specific sequestration factors based on UK soil data from peer-reviewed studies.
	Managed hedgerow planted more than 15 years ago	87, 88, 89	Based on the length and width of managed hedges – gives age-specific sequestration factors based on UK soil data from peer-reviewed studies.
	Large growth with trees	25, 99, & 100	Based on the length and width of large growth hedges with trees – i.e. those trimmed or laid on an irregular basis, forming large structures with in line trees. Sequestration factors based on averages from peer reviewed studies.
Perennial crops	Top fruit, stone fruit and nuts	26	Average sequestration values per hectare. Includes biomass only – soil and grass sequestration excluded.
	Grape vines	28	Covers sequestration in biomass only, not soils.
	Miscanthus	29	Sequestration rates in biomass and soils
	Willow & poplar	30	Covering sequestration in both soils and biomass
Field margins	Uncultivated	25	Area of field margins that are permanently uncultivated. Sequestration rates include soil carbon.
Wetlands	Permanent	13	Area of permanent peaty wetland that is ungrazed

Land use change	(various)	23	Changes which result in losses of carbon, such as woodland to arable. This is from carbon losses in soils and biomass. These are considered to be uncommon in a UK setting, but must be accounted for if they occur. These are one off losses. Users should not enter values in here if they have also calculated SOM measurements for exactly these areas of land use change - though this is thought to be an unlikely occurrence.
	Marshy grassland to degraded wetland	44	Sequestration in biomass and soils on a continuous basis. Users should not enter data here if they have included SOM measurements of the same area.
Habitats/ Higher tier stewardship	(various)	44	Sequestration in biomass and soils on a continuous basis for various habitats, as defined in the Countryside Stewardship Scheme for higher level scheme (HLS) options. The underlying data is based on mid-tier options, and only HLS schemes with an equivalent mid-tier option in the study are included. Users should not enter data here if they have included SOM measurements of the same area.
Cultivated peat soils	Peat soils	21	N ₂ O emissions from cultivated peat soils. Also CO ₂ losses from soils – unless users are able to supply SOM results, in which case only the N ₂ O changes are accounted for. Average values are used from the source.
Uncultivated peatland soils	(various)	82	Emissions from varying states of uncultivated peatland in line with the Peatland Carbon Code.
Countryside Stewardship	(various)	63	Sequestration in biomass and soils on a continuous basis for various habitats, as defined in the Countryside Stewardship Scheme. Users should not enter data here if they have included SOM measurements of the same area.

11.10. Processing

This section calculates carbon emitted as a result of the processing of food and drink including common manufacturing inputs.

Data sources: In this section all of the emission factors are proxy figures, but are all allied to real input quantities, not estimates of items used in a process.

Section	Item	Reference	Notes
Sugar	Cane & Beet	105	GFLI figures for cane sugar production

Fermentation	CO ₂ release	113	Direct CO ₂ released from the fermentation process
Processing products	<i>Various</i>	80	Proxy figures for processing input
	CO ₂ canisters	N/A	Enter the volume of CO ₂ used
	Granulated Sugar	62	Based on cradle to gate for british sugar
Cleaning Products detergents, etc	<i>Various</i>	103	Product specific emissions factors
Packaging	Wine bottles	108	Emissions factor per bottle for 750cl glass wine bottle
	Recycled glass bottle	71	Emissions factor for Encirc recycled "green glass wine bottle" 750cl
	Jars and Bottles	108	Proxy figures for packaging input
	Corks	95	Proxy figures for packaging input
	Crates and Packaging	108	Proxy figures for packaging input
	<i>Various</i>	108	Proxy figures for packaging input
	Refrigeration	Refrigerant usage	12
Water	Mains water	107	Use of mains water
	Mains waste water	107	All waste water released to a mains treatment system
	Non-mains	N/A	Figure simply recorded as water use. No emissions specifically – any fuel or electricity used in pumping or treatment will be picked up under Fuels.

12. Other Calculations we use

Fat and protein corrected milk (FPCM)

To calculate the milk KPI (kg CO₂e per kg FPCM) we use the following equation from the FAO 2010 that corrects to the energy equivalent in milk of 4% fat and 3.3% protein (referenced in 81). If the user does not enter a fat or protein content of their milk, the Calculator assumes 4% fat and 3.2% protein. The calculator also assumes 1 litre = 1.035 kg.

$$\text{kg FPCM}_{[4\%F, 3.2\%CP]} = (0.337 + 0.116 * \text{fat \%} + 0.06 * \text{protein \%})$$



Conversions from individual GHG emissions to CO₂e

The emissions factors for some items in the calculator come from sources such as individual GHG emissions. For example, when accounting for crop residue emissions it is necessary to calculate the direct and indirect N₂O emissions. The calculations provide a value for the quantity of N₂O released, which we then convert into CO₂e per N₂O in accordance with the IPCC guidelines. The three main GHGs are calculated using the following ratios under GWP100 (53):

CO₂ to CO₂e per CO₂ = 1 : 1

CH₄ to CO₂e per CH₄ = 28 : 1

N₂O to CO₂e per N₂O = 265 : 1

13. What farm business information do users enter?

At the start of all reports users are asked to input information about their farm business. Our team is working to ensure we remain best in class for the privacy and data security of your farm business information. Details of how we process your data is outlined in our Terms and Conditions of Software Access and Use, but to summarise this, we do not use or sell this information for purposes other than helping us calculate your carbon footprint and presenting this back to you.

The following farm business information is asked for in reports:

1. **Business category** (s) - Arable, Beef, Dairy, Fruit, Lowland grazing, Mixed (arable/livestock), Other, Pigs, Potatoes, Poultry - layers, Poultry - meat, Sheep, Upland grazing, Upland grazing with common land, Vegetables, Vineyards, Processing, Wineries, Non-agricultural business, Market garden
2. **Farm area** - each of the following categories of land use, in hectares:
 - Cultivated land - all arable and horticulture land that involves soil cultivations (or non/min-till systems)
 - Grassland - temporary and/or permanent grassland, generally used for livestock grazing, and/or forage
 - Non-cropping land - any land not falling under cultivated land or grassland. For example woodland, scrub or other uses which are not generally used for agricultural or horticultural use.
3. **Postcode** - which helps us locate the area where you farm for the purposes of UK benchmarking of results
4. **Certification** - Businesses can mark any certification or assurance schemes they belong to.



- 5. Farm Business identification number** – Businesses can enter an identification number relevant to them, e.g. in England this may be your SFI number. In future upgrades we envisage this number to be used to better connect your report to your farm, and therefore help when integrating outside services like mapping, and other software you may use on your farm.

14. References v.1.6.4 (1 April 2025)

- 1 – SUPERSEDED Department for Business, Energy & Industrial Strategy (2020). 2020 Government greenhouse gas conversion factors for company reporting. Accessed on 16/03/2023
<https://www.gov.uk/government/publications/greenhouse-gas-reporting-conversion-factors-2020>
- 1a – SUPERSEDED Department for Business, Energy & Industrial Strategy (2020). 2020 Government greenhouse gas conversion factors for company reporting: methodology. Accessed on 16/03/2023
https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/901692/conversion-factors-2020-methodology.pdf on 16/03/2023
- 2 – Hammond & Jones (2011). The Inventory of Carbon & Energy (ICE) database v2.0.
- 2a – Jones (2019). The Inventory of Carbon & Energy (ICE) database v3.0. Accessed on 16/03/2023
<https://circularecology.com/embodied-carbon-footprint-database.html>
- 3 – Williams et al. (2006). Determining the environmental burdens and resource use in the production of agricultural and horticultural commodities. DEFRA project report ISO205. Accessed on 16/03/2023
<https://randd.defra.gov.uk/ProjectDetails?ProjectID=11442>
- 4 – SUPERSEDED Brown et al. (2017). UK Greenhouse Gas Inventory, 1990 to 2017: Annual Report for submission under the Framework Convention on Climate Change. Accessed on 20/03/2023 https://naei.beis.gov.uk/reports/reports?report_id=981
- 4a – SUPERSEDED Brown et al. (2017). Annexes to the UK Greenhouse Gas Inventory, 1990 to 2017: Annual Report for submission under the Framework Convention on Climate Change. Accessed on 20/03/2023
https://uk-air.defra.gov.uk/assets/documents/reports/cat09/1905151124_ukghgi-90-17_Annexes_Issue_2_final.pdf
- 5 – Andersen et al. (2010). Quantification of Greenhouse Gas Emissions from Windrow Composting of Garden Waste. Journal of Environmental Quality 39(2): 713–724 <https://doi.org/10.2134/jeq2009.0329>
- 6 – Cuttle et al. (2003) A Review of Leguminous Fertility-Building Crops, with Particular Reference to Nitrogen Fixation and Utilisation Written as a Part of Defra Project OF0316 “The Development of Improved Guidance on the Use of Fertility-Building Crops in Organic Farming”. Institute of Grassland and Environmental Research: Aberystwyth, Wales, 2003.
- 7 – Phong (2012). Greenhouse Gas Emissions from Composting and Anaerobic Digestion Plants. INRES, Institute of Crop Science and Resource Conservation. Bonn, D-53115.
- 8 – Amon et al. (1999). Emissions of NH₃, N₂O and CH₄ from composted and anaerobically stored farm yard manure. Ramiran 98 posters presentations. Accessed on 16/03/2023 <http://ramiran.uvlf.sk/doc98/FIN-POST/AMON-BAR.pdf>
- 9 – SUPERSEDED FCT calculations
- 10 – SUPERSEDED Woodland Carbon Code. (2018). Carbon Lookup tables v2.0. Accessed on 30/05/2022
<https://www.woodlandcarboncode.org.uk/news/version-2-0-of-the-wcc-launched?highlight=WyJsb29rdXAiXQ==>
- 11 – Clark (2007). Cover crops—United States—Handbooks, manuals, etc. Sustainable Agriculture Network. 3rd edition.
- 12 – GHG protocol (2017). Calculating HFC and PFC emissions from the manufacturing, servicing, and/or disposal of refrigeration and air-conditioning equipment. Calculation worksheets v1.0. Accessed on 30/05/2022
https://view.officeapps.live.com/op/view.aspx?src=https%3A%2F%2Fghgprotocol.org%2Fsites%2Fdefault%2Ffiles%2Fhfc-pfc_0.xls
- 13 – Taylor et al. (2010). Measuring holistic carbon footprints for lamb and beef farms in the cambrian mountains initiative. CCW Policy Research Report No. 10/8.
- 14 – Bentrup et al. (2016). Carbon footprint analysis of mineral fertilizer production in Europe and other world regions. Conference paper. Accessed on 30/05/2022
https://www.researchgate.net/publication/312553933_Carbon_footprint_analysis_of_mineral_fertilizer_production_in_Europe_and_other_world_regions
- 15 – SUPERSEDED Berners-Lee (2010). How bad are bananas? The carbon footprint of everything. Profile Books, London



- 16 – Warwick HRI (2009). Preliminary assessment of greenhouse gases associated with growing media materials. DEFRA project report IF0154 <http://randd.defra.gov.uk/Default.aspx?Module=More&Location=None&ProjectID=15967>
- 17 – Wiltshire et al. (2008). Scenario building to test and inform the development of a BSI method for assessing greenhouse gas emissions from food (Technical annex to the final report). DEFRA project report FO0404 submitted by ADAS. Accessed 02/05/2023
<https://repository.rothamsted.ac.uk/item/8q33x/scenario-building-to-test-and-inform-the-development-of-a-bsi-method-for-assessing-greenhouse-gas-emissions-from-food-technical-annex-to-final-report-on-defra-project-no-fo0404> *
- 18 – SUPERSEDED GFLI (2020). Database of livestock feeds and environmental impacts. Accessed 30/05/2022
<http://globalfeedlca.org/gfli-database/gfli-database-tool/>
- 19 – SUPERSEDED FCT calculations
- 20 – SUPERSEDED Correspondence with David McNaughton (Soya UK Managing Director) on crop yields and residues
- 21 – Taft et al. (2017) GHG from intensively managed peat soils in an arable production system. *Agriculture, Ecosystems & Environment*. 237: 162-172.
- 22 – Axe et al. (2017) Carbon storage in hedge biomass – A case study of actively managed hedges in England. *Agriculture, Ecosystems & Environment*. 250: 81-88. <https://doi.org/10.1016/j.agee.2017.08.008>
- 23 – Ostle et al. (2009). UK land use and carbon sequestration. *Land Use Policy* 26S: S274-S283.
<https://doi.org/10.1016/j.landusepol.2009.08.006>
- 24 – Chishna et al (2010) Embodied carbon in natural building stone in Scotland. Historic Scotland, Technical Conservation Group. Technical Paper 7. SiSTech Ltd and Harold-Watt University.
- 25 – Falloon et al (2004) Managing field margins for biodiversity and carbon sequestration: A Great Britain case study. *Soil Use and Management*. 20 (2): 240-247.
- 26 – Kerckhoffs and Reid (2007). Carbon sequestration in the standing biomass of orchard crops in New Zealand. NZ Institute for Crop & Food Research Ltd. report for Horticulture New Zealand Ltd.
- 27 – Carlisle et al. (2010). California vineyard greenhouse gas emissions: assessment of the available literature and determination of research needs. California sustainable wine growing Alliance. Accessed on 30/05/2022
https://www.sustainablewinegrowing.org/docs/CSWA%20GHG%20Report_Final.pdf
- 28 – Vicente-Vicente et al. (2016) Soil carbon sequestration rates under Mediterranean woody crops using recommended management practices: A meta-analysis. *Agriculture, Ecosystems & Environment*. 235: 204-214.
- 29 – Dondini et al. (2009). The potential of Miscanthus to sequester carbon in soils: comparing field measurements in Carlow, Ireland to model predictions. *GCB Bioenergy* 1: 413-425. <https://doi.org/10.1111/j.1757-1707.2010.01033.x>
- 30 – Rytter (2012) The potential of willow and poplar plantations as carbon sinks in Sweden. *Biomass and Bioenergy*. 36:86-95. <https://doi.org/10.1016/j.biombioe.2011.10.012>
- 31 – Grogan and Matthews (2002). A modelling analysis of the potential for soil carbon sequestration under short rotation coppice willow bioenergy plantations. *Soil Use and Management* 18: 175-183. <https://doi.org/10.1111/j.1475-2743.2002.tb00237.x>
- 32 – Ventura et al (2019) Carbon balance and soil carbon input in a poplar short rotation coppice plantation as affected by nitrogen and wood ash application. *New Forests*. 50. 969-990.
- 33 – Turner et al (2015) Greenhouse gas emission factors for recycling of source-segregated waste materials. *Resources, Conservation and Recycling*. 105 (A): 186-197.
- 34 – SUPERSEDED Personal communications with Chris Foss (Wine GB)
- 35 – COFALEC (2015). Carbon footprint of yeast produced in the European Union. Produced by PriceWaterhouseCooper for COFALEC. Accessed 30/05/2022
https://cofalec.com/wp-content/uploads/2022/03/20120327155707_Yeast_Carbon_Footprint_COFALEC_28english-version29.pdf
- 36 – Nica and Woinarocschy (2010) Environmental Assessment of Citric Acid production. *UPB Scientific Bulletin, Series B. Chemistry and Materials Science*. 72 (3):45-56.
- 37 – AHDB & HGCA (2014). Carbon footprint decision tool. 10. Field Operations. Accessed 21/03/2023
[https://view.officeapps.live.com/op/view.aspx?src=https%3A%2F%2Fprojectblue.blob.core.windows.net%2Fmedia%2FDefault%2FTools%2FTool%2520Download%2FAHDB%2520carbon%2520footprinting%2520tool%2520\(2014\).xism&wdOrigin=BROWSELINK](https://view.officeapps.live.com/op/view.aspx?src=https%3A%2F%2Fprojectblue.blob.core.windows.net%2Fmedia%2FDefault%2FTools%2FTool%2520Download%2FAHDB%2520carbon%2520footprinting%2520tool%2520(2014).xism&wdOrigin=BROWSELINK)
- 38 – Moller et al. (2009) Anaerobic digestion and digestate use: accounting of greenhouse gases and global warming contribution. *Waste Manag Res*. 27 (8): 813-24.



- 39 - Vergana & Silver (2019) GHG emissions from windrow composting of organic wastes: Patterns and emissions factors. *Environmental Research Letters*. 14 (12) 124027.
- 40 - Audsley et al. (2009) Estimation of the greenhouse gas emissions from agricultural pesticide manufacture and use. Cranfield University. 10. Accessed 30/05/2022
https://dspace.lib.cranfield.ac.uk/bitstream/handle/1826/3913/Estimation_of_the_greenhouse_gas_emissions_from_agricultural_pesticide_manufacture_and_use%E2%80%90902009.pdf?sequence=1
- 41 - Yara (2017). Yara International ASA. Carbon footprint - fertilizer products. Verified by DNV GL. Accessed on 25/04/2023
https://www.yara.co.uk/contentassets/a6e77004605040aea339577f909d5368/yara-carbon-footprint_verification_statement.pdf
- 42 - CF Fertiliser range (under reconsideration, reference material unavailable)
- 43 - Schwarzbeck et al (2015) Determining national greenhouse gas emissions from waste-to-energy using the Balance Method Determining national greenhouse gas emissions from waste-to-energy using the Balance Method. *Waste Management*. 49:263-271. <https://doi.org/10.1016/j.wasman.2016.01.025>
- 44 - Warner et al. (2020b). Establishing a field-based evidence base for the impact of agri-environment options on soil carbon and climate change mitigation – phase 2. Final Report. Work package number: ECM50416. Evidence Programme Reference number: RP04176. Natural England.
- 45 - Farm Carbon Toolkit: Soil Carbon Project (ongoing). See <https://farmcarbontoolkit.org.uk/soil-carbon-project/> for more information.
- 46 - Personal communications with Joseph Barnes (Saria UK)
- 47 - Fertilizers Europe (2011). Carbon footprint reference values - mineral fertilizer carbon footprint reference values: 2011.
- 48 - Brenttrup et al (2018) Updated carbon footprint values for mineral fertilizer from different world regions. LCA Food 2018 and LCA AgriFood Asia 2018: (1-B) From Farm to Table. Conference paper accessed on 30/05/2022
https://www.researchgate.net/publication/329774170_Updated_carbon_footprint_values_for_mineral_fertilizer_from_different_world_regions
- 49 - Sylvester-Bradley et al. (2015). Minimising nitrous oxide intensities of arable crop products (MIN-NO). AHDB Cereals & Oilseeds/ Project Report No. 548. Accessed on 30/05/2022
<https://projectblue.blob.core.windows.net/media/Default/Research%20Papers/Cereals%20and%20Oilseed/pr548-abstract-and-executive-summary.pdf>
- 50 - AHDB (2017). Nutrient Management Guide - RB209. Accessed on 30/05/2022 <https://ahdb.org.uk/RB209>
- 51 - Thorman et al (2020) Towards Country-Specific Nitrous Oxide Emission Factors for Manures Applied to Arable and Grassland Soils in the UK. *Frontiers in Sustainable Food Systems*. 4:62.
- 52 - Liang & Kasimir (2019) Chapter 11: N₂O Emissions from Managed Soils, and CO₂ Emissions from Lime and Urea Application. Refinement to 2006 IPCC Guidelines for National Greenhouse Gas Inventories (pp. 11.1-11.48) Publisher: Intergovernmental Panel on Climate Change.
- 53 - IPCC (2020). Climate Change and Land - An IPCC Special Report on climate change, desertification, land degradation, sustainable land management, food security, and greenhouse gas fluxes in terrestrial ecosystems. Summary for policy makers. ISBN 978-92-9169-154-8. Available at <https://www.ipcc.ch/srccl/chapter/summary-for-policymakers/>
- 54 - Haverkort and Hillier (2011). Cool Farm Tool - Potato: Model Description and Performance of Four Production Systems. *Potato Res.* 54, 355-369 <https://doi.org/10.1007/s11540-011-9194-1>
- 55 - SUPERSEDED Department for Business, Energy & Industrial Strategy (2021). UK Government GHG Conversion Factors for Company Reporting 2021. Accessed on 30/05/2021
<https://www.gov.uk/government/publications/greenhouse-gas-reporting-conversion-factors-2021>
- 56 - PET Recycling Team website (2017). Certificate of carbon footprint for PCF Model ALPHA Bottles rPET produced using EcoInvent 3.3. Accessed on 30/05/2021 <https://petrecyclingteam.com/en/excellent-co2-balance>
- 57 - SUPERSEDED Idemat database (2020). ECO-costs 2017 v1.6. Accessed on 30/05/2021
<https://view.officeapps.live.com/op/view.aspx?src=https%3A%2F%2Fwww.ecocostsvalue.com%2FEVR%2Fimg%2FIdematapp2020.xlsx&wdOrigin=BROWSELINK>
- 58 - SUPERSEDED West (2021). Woodland Carbon Code Carbon Calculations Spreadsheet Version 2.4. Accessed 30/05/2021
https://www.woodlandcarboncode.org.uk/images/Spreadsheets/WCC_CarbonCalculationSpreadsheet_Version2.4_March2021.xlsx



- 59 – SUPERSEDED Brown et al. (2021). UK Greenhouse Gas Inventory 1990 to 2019: Annual Report for submission under the Framework Convention on Climate Change. Department for Business, Energy & Industrial Strategy. Accessed on 30/05/2022 https://uk-air.defra.gov.uk/assets/documents/reports/cat09/2105061125_ukghgi-90-19_Main_Issue_1.pdf
- 59a – SUPERSEDED Brown et al. (2021). Annexes to the UK Greenhouse Gas Inventory 1990 to 2019: Annual Report for submission under the Framework Convention on Climate Change. Department for Business, Energy & Industrial Strategy. Accessed on 30/05/2022 https://uk-air.defra.gov.uk/assets/documents/reports/cat09/2106091119_ukghgi-90-19_Annex_Issue_2.pdf
- 60 – Bizarro et al. (2021). Potential carbon footprint reduction for reclaimed asphalt pavement innovations. Sustainability 13(3):1382 <https://doi.org/10.3390/su13031382>
- 61 – GHG Protocol (2014). Agricultural Guidance Interpreting the Corporate Accounting and Reporting Standard for the agricultural sector. GHG Protocol Agricultural Guidance. Accessed on 02/03/23 https://ghgprotocol.org/sites/default/files/standards/GHG%20Protocol%20Agricultural%20Guidance%20%28April%2026%29_0.pdf
- 62 – Carbon Trust (2021). Certification Letter – British Sugar – 2020 LimeX extension. Carbon Trust CERT-10235
- 63 – Warner et al. (2020a). Establishing a field-based evidence base for the impact of agri-environment options on soil carbon and climate change mitigation – phase 1. Final Report. Work package number: ECM50416. Evidence Programme Reference number: RP04176. Natural England.
- 64 – SUPERSEDED Department for Business, Energy & Industrial Strategy (2022) Greenhouse gas reporting: conversion factors 2022. Accessed on 04/01/2023 <https://www.gov.uk/government/publications/greenhouse-gas-reporting-conversion-factors-2022>
- 65 – SUPERSEDED Brown et al. (2022) UK Greenhouse Gas Inventory, 1990 to 2020. Department for Business, Energy & Industrial Strategy. Accessed on 05/01/2023 https://uk-air.defra.gov.uk/assets/documents/reports/cat09/2206220830_ukghgi-90-20_Main_Issue1.pdf
- 66 – SUPERSEDED Brown et al. (2022) UK Greenhouse Gas Inventory 2020 annexes. Department for Business, Energy & Industrial Strategy. Accessed 05/01/2023 https://naei.beis.gov.uk/reports/reports?report_id=1072
- 67 – Wilms et al. (2022). Macronutrient profile in milk replacer or a whole milk powder modulates growth performance, feeding behavior, and blood metabolites in ad libitum-fed calves. J. Dairy Sci. 105:6670–6692 <https://doi.org/10.3168/jds.2022-21870>
- 68 – Finnegan et al. (2017). Environmental impacts of milk powder and butter manufactured in the Republic of Ireland. Science of the Total Environment 579 (2017) 159–168 <http://dx.doi.org/10.1016/j.scitotenv.2016.10.237>
- 69 – Sánchez et al. (2012). Comparison of Life Cycle energy consumption and GHG emissions of natural gas, biodiesel and diesel buses of the Madrid transportation system. Energy 47(1):174–198 <https://doi.org/10.1016/j.energy.2012.09.052>
- 70 – Smyth et al. (2015) Developing Peatland Carbon Metrics and Financial Modelling to Inform the Pilot Phase UK Peatland Code. Report to Defra for Project NR0165, Crichton Carbon Centre, Dumfries.
- 71 – Carbon Intelligence (2021) Encirc LCA for wine bottle, green glass, conducted by Carbon Intelligence.
- 72 – Budsberg et al. (2020). Production routes to bio-acetic acid: life cycle assessment. Biotechnol Biofuels 13:154 <https://doi.org/10.1186/s13068-020-01784-y>
- 73 – Bellboom et al. (2015). Environmental impacts of phosphoric acid production using di-hemihydrate process: a Belgian case study. Journal of Cleaner Production 108A: 978–986 <https://doi.org/10.1016/j.jclepro.2015.06.141>
- 74 – Naukkarinen (2023). Life Cycle Assessment Study of a Sulfuric Acid Manufacturing Process in the Chemi-cal Pulping Industry. Masters thesis, Lappeenranta–Lahti University of Technology LUT. Accessed 27/04/2023 https://lutpub.lut.fi/bitstream/handle/10024/165170/Thesis_Naukkarinen_Martta.pdf?sequence=1
- 75 – Origin (2020). RSK ADAS Limited certificate of cradle-to-gate carbon footprint at the plant gate (Origin Newport) of Origin CAN
- 76 – Origin (2020). RSK ADAS Limited certificate of cradle-to-gate carbon footprint at the plant gate (Origin Newport) of Origin 14-14-21 + 7SO₃ + 0.02B
- 77 – Origin (2020). RSK ADAS Limited certificate of cradle-to-gate carbon footprint at the plant gate (Origin Newport) of Origin 16-16-16 + 7SO₃ + 0.02B
- 78 – Origin (2020). RSK ADAS Limited certificate of cradle-to-gate carbon footprint at the plant gate (Origin Newport) of Origin 10-10-20 + 7SO₃ + 0.02B
- 79 – Ogle et al. (2019). Refinement to 2006 IPCC Guidelines for National Greenhouse Gas Inventories. Volume 4 – Agriculture, forestry and other land use. Chapter 2 – Generic methodologies applicable to multiple land use categories (pp. 2.33) Publisher:



Intergovernmental Panel on Climate Change.

https://www.ipcc-nggip.iges.or.jp/public/2019rf/pdf/4_Volume4/19R_V4_Ch02_Generic%20Methods.pdf

80 - International Organisation of Vine and Wine (2017). Methodological recommendations for accounting for the GHG balance in the vitiviculture sector. Paris, France. ISBN 979-10-91799-75-1

81 - Baldini et al. (2017). A critical review of the recent evolution of Life Cycle Assessment applied to milk production. *Journal of Cleaner Production* 140: 421e435 <http://dx.doi.org/10.1016/j.jclepro.2016.06.078>

82 - Evans et al. (2022). Aligning the Peatland Code with the UK Peatland Inventory. Report to Defra and the IUCN Peatland Programme, March 2022 (Updated January 2023)

83 - Farm Carbon Toolkit (2023). Report for Mercian Seed potato supplier farm (England) for cropping year 2022.

[https://calculator.farmcarbontoolkit.org.uk/sites/default/files/83.%20FCC%20Report%20\(2022\)%20Mercian%20English%20Seed%20Potatoes.pdf](https://calculator.farmcarbontoolkit.org.uk/sites/default/files/83.%20FCC%20Report%20(2022)%20Mercian%20English%20Seed%20Potatoes.pdf)

84 - Farm Carbon Toolkit (2023). Report for Mercian Seed potato supplier farm (Scotland) for cropping year 2022.

[https://calculator.farmcarbontoolkit.org.uk/sites/default/files/84.%20FCC%20Report%20\(2022\)%20Mercian%20Scottish%20seed%20potatoes.pdf](https://calculator.farmcarbontoolkit.org.uk/sites/default/files/84.%20FCC%20Report%20(2022)%20Mercian%20Scottish%20seed%20potatoes.pdf)

85 - SUPERSEDED Department for Energy Security and Net Zero (2023). Greenhouse Gas Reporting: Conversion Factors 2023 (flat file .csv) Published 07/06/2023, accessed on 01/12/2023

<https://www.gov.uk/government/publications/greenhouse-gas-reporting-conversion-factors-2023>

86 - SUPERSEDED Department for Energy Security and Net Zero (2023). Greenhouse Gas Reporting: Conversion Factors 2023 (full file .xls) Published 07/06/2023, accessed on 01/12/2023

<https://www.gov.uk/government/publications/greenhouse-gas-reporting-conversion-factors-2023>

87 - Drexler, S., Thiessen, E., & Don, A. (2023). Carbon storage in old hedgerows: The importance of below-ground biomass. *GCB Bioenergy*, 16, e13112. <https://doi.org/10.1111/gcbb.13112>

88 - Biffi, S., Chapman, P., Grayson, R.P., Ziv, G. (2022). Soil carbon sequestration potential of planting hedgerows in agricultural landscapes. *Journal of Environmental Management*, 307, 114484. <https://doi.org/10.1016/j.jenvman.2022.114484>

89 - Biffi, S., Chapman, P., Grayson, R.P., Ziv, G. (2023). Planting hedgerows: Biomass carbon sequestration and contribution towards net-zero targets. *Science of the Total Environment*, 892, 164482. <https://doi.org/10.1016/j.scitotenv.2023.164482>

90 - Wang, Z., Chen, J., Mao, S., Han, Y., Chen, F., Zhang, L., Li, Y., & Li, C., (2017) Comparison of greenhouse gas emissions of chemical fertilizer types in China's crop production. *Journal of Cleaner Production*. 141, 1267-1274.

91 - Meinrenken, Christoph J; Chen, Daniel; Esparza, Ricardo A; Iyer, Venkat; Prasad, Aruna; Paridis, Sally; et al. (2022). The Carbon Catalogue public database – Carbon footprints of 866 commercial products across 8 industry sectors and 5 continents. figshare. Dataset. <https://doi.org/10.6084/m9.figshare.16908979.v1>

92a - SUPERSEDED Brown et al. (2023) UK Greenhouse Gas Inventory, 1990 to 2021. Department for Energy Security and Net Zero. Accessed on 08/03/2024

https://uk-air.defra.gov.uk/assets/documents/reports/cat09/2304171441_ukghgi-90-21_Main_Issue1.pdf

92b - SUPERSEDED Brown et al. (2023) UK Greenhouse Gas Inventory, 1990 to 2021 Annexes. Department for Energy Security and Net Zero. Accessed on 08/03/2024

https://uk-air.defra.gov.uk/assets/documents/reports/cat09/2304171442_ukghgi-90-21_Annex_Issue1.pdf

92c - SUPERSEDED Brown et al. (2023) UK Greenhouse Gas Inventory, 1990 to 2021 Supplementary file with emission factors for the agriculture sector. Department for Energy Security and Net Zero. Accessed on 08/03/2024

https://view.officeapps.live.com/op/view.aspx?src=https%3A%2F%2Fuk-air.defra.gov.uk%2Fassets%2Fdocuments%2Freports%2Fcat09%2F2304171445_Supplementary_file_EFs_UK_inventory_agriculture_2023.xlsx&wdOrigin=BROWSELINK

93 - Gavrilova et al. (2019) Chapter 10: Emissions from Livestock and Manure Management. Refinement to 2006 IPCC Guidelines for National Greenhouse Gas Inventories (pp. 10.1 – 10.207) Publisher: Intergovernmental Panel on Climate Change.

94 - Liang & Kasimir (2019) Chapter 11: N2O Emissions from Managed Soils, and CO2 Emissions from Lime and Urea Application. Refinement to 2006 IPCC Guidelines for National Greenhouse Gas Inventories (pp. 11.1-11.48) Publisher: Intergovernmental Panel on Climate Change. https://www.ipcc-nggip.iges.or.jp/public/2019rf/pdf/4_Volume4/19R_V4_Ch11_Soils_N2O_CO2.pdf

95 - Vogtlander (2024). Idemat dataset V1-2. Accessed on 05/03/2024 [Idemat_2024-V2-1.xlsx \(live.com\)](https://www.live.com/Idemat_2024-V2-1.xlsx)

96a - RB209 (2023). Nutrient Management Guide: Section 1. Principles of Nutrient Management and Fertiliser Use. Accessed on 08/03/2024 <https://ahdb.org.uk/knowledge-library/rb209-section-1-principles-of-nutrient-management-and-fertiliser-use>



- 96b - RB209 (2023). Nutrient Management Guide: Section 2. Organic Materials. Accessed on 08/03/2024
<https://ahdb.org.uk/knowledge-library/rb209-section-2-organic-materials>
- 97 - Steel Insight (2011). Last accessed on 25/03/2024
<https://www.building.co.uk/home/steel-insight-structural-steelwork/5026908.article>
- 98 - ForFarmers (2024) Data on feed composition provided by ForFarmers, December 2023.
- 99 - Crossland (2015). The carbon sequestration potential of hedges managed for woodfuel. The Organic Research Centre. Last accessed on 25/03/2024
https://www.organicresearchcentre.com/manage/authincludes/article_uploads/project_outputs/TWECOM%20ORC%20Carbon%20report%20v1.0.pdf
- 100 - Taylor et al. (2010). Measuring holistic carbon footprints for beef and lamb in the Cambrian Mountains Initiative. CCW Policy Research Report No. 10/8
- 101 - Robertson et al. (2012). Economic, biodiversity, resource protection and social values of orchards: A study of six orchards by the Herefordshire Orchards Community Evaluation Project. Natural England Commissioned Report NECR090
- 102 - SUPERSEDED Department for Energy Security and Net Zero (2023). Greenhouse Gas Reporting: Conversion Factors 2023 (full file .xls) Published 07/06/2023, accessed on 01/12/2023
<https://www.gov.uk/government/publications/greenhouse-gas-reporting-conversion-factors-2023>
- 103 - Evans Vanodine (2024) Technical Hub - Carbon Calculator. Available at:
<https://www.evansvanodine.co.uk/carbon-calculator> Last accessed on 07/10/2024
- 104 - The Woodland Carbon Code (2024) Calculation Spreadsheet. Available at: <https://www.woodlandcarboncode.org.uk/>
- 105 - The Global Feed LCA Institute (2023). Database of livestock feeds and environmental impacts. Available at:
<https://globalfeedlca.org/>
- 106 - International Standard ISO146064-1 (2018) Second Edition 2018-12. Part 1: Specification with guidance at the organization level for quantification and reporting of greenhouse gas emissions and removals. Available at:
<https://www.iso.org/standard/66453.html>
- 107a - Department for Energy Security and Net Zero (2024). Greenhouse Gas Reporting: Conversion Factors 2024 (full file .csv) Published 08/06/2024, accessed on 14/03/2025
<https://www.gov.uk/government/publications/greenhouse-gas-reporting-conversion-factors-2024>
- 107b - Department for Energy Security and Net Zero (2024). Greenhouse Gas Reporting: Conversion Factors 2024 (flat file .csv) Published 08/06/2024, accessed on 14/03/2025
<https://www.gov.uk/government/publications/greenhouse-gas-reporting-conversion-factors-2024>
- 108 - Circular Ecology (2024). The Inventory of Carbon and Energy (ICE) Database Advanced V4.0. Published Dec 2024, accessed on 14/03/2024 <https://circularecology.com/embodied-carbon-footprint-database.html>
- 109 - Idemat database (2024). Data on Eco-costs 2024 V1.0 Accessed on 25/03/2025
- 110 - Marshalls Plc (2023) Environmental Product Declaration Concrete Paving Flags - Standard Wet Pressed.
- 111a - Brown et al. (2024). UK Greenhouse Gas Inventory, 1990 to 2022. Annual Report for Submission under the Framework Convention on Climate Change. Accessed on 25/03/2025. Available at: https://uk-air.defra.gov.uk/library/reports?report_id=1
- 111b - Brown et al. (2024). UK Greenhouse Gas Inventory, 1990 to 2022 Annexes. Accessed on 25/03/2025. Available at: https://uk-air.defra.gov.uk/library/reports?report_id=1136
- 111c - Brown et al. (2024). UK Greenhouse Gas Inventory, 1990 to 2022 Supplementary Information. Accessed on 25/03/2025. Available at: https://uk-air.defra.gov.uk/library/reports?report_id=1137
- 112 - Hutchinsons (2023) Fertiliser Formulations provided by Hutchinsons (2023).
- 113 - Prusova et al. (2023). Capture of fermentation gas from fermentation of grape must. Foods, 12(3), p.574.
- 114 - Webb et al (2024) The impacts of manure application methods on emissions of ammonia, nitrous oxide and on crop response—A review, Agriculture, Ecosystems and Environment 137 (2010) 39–46

END OF LIST



15. Contacting us

We welcome Calculator users to contact the Calculator team with questions, suggestions and comments at any time.

For general enquiries, please email: calculator@farmcarbontoolkit.org.uk or reach out to a member of our team.



Calculator Manager:	Lizzy Parker
Calculator Development officer:	James Pitman
Calculator Development officer:	Grace Wardell
Data Scientist:	Izzy Peters
Data Assistant:	Calum Adams
Customer Services Officer:	Michael Brown (contact)

16. Copyright and use

This document is subject to copyright © Farm Carbon Toolkit, 2025. We would suggest you share this with your team or point other users to this document where you think they would benefit from it. In case it changes - send them the link to our [resources page](#) so they can see the latest version.